HEDIS® Tip Sheet



Learn How to Improve Your HEDIS¹ Rate for Osteoporosis Screening

Use this tip sheet to review key details of the Osteoporosis Screening in Older Women (OSW) measure, best practices and resources.

Measure	 Women 65–75 years of age who received osteoporosis screening. One or more osteoporosis screening tests on or between the member's 65th birthday and December 31 of the measurement year. 	
Exclusions	 Patients who meet the following criteria: Had a claim/encounter for osteoporosis therapy any time in the member's history through December 31 of the year prior to the measurement year. Received a dispensed prescription to treat osteoporosis anytime on or between January 1 three years prior to the measurement year through December 31 of the year prior to the measurement year. Hospice or using hospice services anytime during the measurement year. Palliative care during the measurement year. Members 66 and older as of December 31 of the measurement year enrolled in an Institutional Special Needs Plan (I-SNP) or living long-term in an institution any time during the measurement year. 	
	measurement year with frailty and advanced illness.	



Often called a silent disease, osteoporosis reduces bone mass through structural deterioration, resulting in compromised bone strength. Osteoporosis is more common in women than in men. One in two women will have an osteoporosis-related fracture in their lifetime.² Routine screenings can help identify osteoporosis and allow members to take preventive measures.

(continued)

	Submit a claim for osteoporosis screening.Set up a process for bone mineral density (BMD) testing:	
	 To determine if a member had a BMD test, query enrolled members in the age span noted. 	
	 Provide a BMD prescription and urge patients to complete the screening soon 	
	 Make sure the screening site notifies you with results in a timely manner. 	
Best	 Set up a follow-up visit to discuss the results at the patient's next visit. 	
practices	• Evaluate women for risk factors that would increase the risk of osteoporosis. Some risk factors include low body weight, excessive alcohol intake, current tobacco use, history of fractures, and use of certain medications.	
	• Discuss osteoporosis prevention methods with your patients, such as calcium, vitamin D supplements, and weight-bearing exercises.	
	• Have culturally appropriate, tailored educational messaging that addresses the risk and importance of osteoporosis in racial and ethnic minority populations that have lower screening rates.	

		Use the appropriate service codes when billing for OSW screenings CPT Copyright 2017 American Medical Association. All rights reserved. CPT [®] is a registered trademark of the American Medical Association.			
C	Codes	Tests/medications	CPT codes	HCPCS codes	
		Osteoporosis screening tests	76977, 77078, 77080, 77081, 77085		
		Osteoporosis medications		J0897, J1740, J3110, J3111, J3489	
		Long-acting osteoporosis medications during an inpatient stay		J0897, J1740, J3489	

Resources

[•] Final Recommendation Statement, Osteoporosis to Prevent Fractures: Screening. U.S. Preventive Services Task Force at

www.uspreventives ervices task force.org/uspstf/recommendation/osteoporos is-screening.

[•] National Committee for Quality Assurance (NCQA), HEDIS Measurement Year 2022, Volume 2: Technical Specifications for Health Plans, 2021.