

Clinical Policy: Ravulizumab-cwvz (Ultomiris)

Reference Number: CP.PHAR.415

Effective Date: 06.01.19

Last Review Date: 12.20

Line of Business: Commercial, TBD HIM*, Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

***For Health Insurance Marketplace members**, if request is through the pharmacy benefit, this policy applies only when the referenced drug is on the health plan approved formulary. Request for non-formulary drugs must be reviewed using the policy: HIM.PA.103.

Description

Ravulizumab-cwvz (Ultomiris[®]) is a complement inhibitor.

FDA Approved Indication(s)

Ultomiris is indicated for the treatment of:

- Adult patients with paroxysmal nocturnal hemoglobinuria (PNH)
- Adult and pediatric patients one month of age and older with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy (TMA)

Limitation(s) of use: Ultomiris is not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Ultomiris is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Paroxysmal Nocturnal Hemoglobinuria** (must meet all):

1. Diagnosis of PNH;
2. Prescribed by or in consultation with a hematologist;
3. Age \geq 18 years;
4. Flow cytometry shows detectable GPI-deficient hematopoietic clones or \geq 5% PNH cells;
5. Member meets one of the following (a or b):
 - a. History of \geq 1 red blood cell transfusion in the past 24 months and (i or ii):
 - i. Documentation of hemoglobin $<$ 7 g/dL in members without anemia symptoms;
 - ii. Documentation of hemoglobin $<$ 9 g/dL in members with anemia symptoms;
 - b. History of thrombosis;
6. Dose does not exceed the following (a, b, and c):

- a. Loading dose on Day 1 (i, ii, or iii):
 - i. Weight \geq 40 to $<$ 60 kg: 2,400 mg;
 - ii. Weight \geq 60 to $<$ 100 kg: 2,700 mg;
 - iii. Weight \geq 100 kg: 3,000 mg;
- b. If member is switching therapy from Soliris[®], administration of the loading dose should occur 2 weeks after the last Soliris infusion;
- c. Maintenance dose on Day 15 and every 8 weeks thereafter (i, ii, or iii):
 - i. Weight \geq 40 to $<$ 60 kg: 3,000 mg;
 - ii. Weight \geq 60 to $<$ 100 kg: 3,300 mg;
 - iii. Weight \geq 100 kg: 3,600 mg.

Approval duration: 6 months

B. Atypical Hemolytic Uremic Syndrome (must meet all):

1. Diagnosis of aHUS (i.e., complement-mediated HUS);
2. Prescribed by or in consultation with a hematologist or nephrologist;
3. Age \geq 1 month;
4. Member has signs of TMA as evidenced by all of the following (a, b, and c):
 - a. Platelet count \leq $150 \times 10^9/L$;
 - b. Hemolysis such as an elevation in serum lactate dehydrogenase (LDH);
 - c. Serum creatinine above the upper limits of normal or member requires dialysis;
5. Documentation that member does not have either of the following:
 - a. A disintegrin and metalloproteinase with thrombospondin type 1 motif, member 13 (ADAMTS13) deficiency;
 - b. STEC-HUS;
6. Dose does not exceed the following (a, b, and c):
 - a. Loading dose on Day 1:
 - i. Weight \geq 5 to $<$ 10 kg: 600 mg;
 - ii. Weight \geq 10 to $<$ 20 kg: 600 mg;
 - iii. Weight \geq 20 to $<$ 30 kg: 900 mg;
 - iv. Weight \geq 30 to $<$ 40 kg: 1,200 mg;
 - v. Weight \geq 40 to $<$ 60 kg: 2,400 mg;
 - vi. Weight \geq 60 to $<$ 100 kg: 2,700 mg;
 - vii. Weight \geq 100 kg: 3,000 mg;
 - b. If member is switching therapy from Soliris, administration of the loading dose should occur 2 weeks after the last Soliris infusion;
 - c. Maintenance dose on Day 15 and at the specified frequency thereafter:
 - i. Weight \geq 5 to $<$ 10 kg: 300 mg every 4 weeks;
 - ii. Weight \geq 10 to $<$ 20 kg: 600 mg every 4 weeks;
 - iii. Weight \geq 20 to $<$ 30 kg: 2,100 mg every 8 weeks;
 - iv. Weight \geq 30 to $<$ 40 kg: 2,700 mg every 8 weeks;
 - v. Weight \geq 40 to $<$ 60 kg: 3,000 mg every 8 weeks;
 - vi. Weight \geq 60 to $<$ 100 kg: 3,300 mg every 8 weeks;
 - vii. Weight \geq 100 kg: 3,600 mg every 8 weeks.

Approval duration: 6 months

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters (a or b):
 - a. PNH:
 - i. Improved measures of intravascular hemolysis (e.g., normalization of LDH);
 - ii. Reduced need for red blood cell transfusions;
 - iii. Increased or stabilization of hemoglobin levels;
 - iv. Less fatigue;
 - v. Improved health-related quality of life;
 - vi. Fewer thrombotic events;
 - b. aHUS:
 - i. Improved measures of intravascular hemolysis (e.g., normalization of LDH);
 - ii. Increased or stabilized platelet counts;
 - iii. Improved or stabilized serum creatinine or estimated glomerular filtration rate (eGFR);
 - iv. Reduced need for dialysis;
3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
 - a. PNH:
 - i. Weight \geq 40 to $<$ 60 kg: 3,000 mg every 8 weeks;
 - ii. Weight \geq 60 to $<$ 100 kg: 3,300 mg every 8 weeks;
 - iii. Weight \geq 100 kg: 3,600 mg every 8 weeks;
 - b. aHUS:
 - i. Weight \geq 5 to $<$ 10 kg: 300 mg every 4 weeks;
 - ii. Weight \geq 10 to $<$ 20 kg: 600 mg every 4 weeks;
 - iii. Weight \geq 20 to $<$ 30 kg: 2,100 mg every 8 weeks;
 - iv. Weight \geq 30 to $<$ 40 kg: 2,700 mg every 8 weeks;
 - v. Weight \geq 40 to $<$ 60 kg: 3,000 mg every 8 weeks;
 - vi. Weight \geq 60 to $<$ 100 kg: 3,300 mg every 8 weeks;
 - vii. Weight \geq 100 kg: 3,600 mg every 8 weeks.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ADAMTS13: a disintegrin and metalloproteinase with thrombospondin type 1 motif, member 13

aHUS: atypical hemolytic uremic syndrome

FDA: Food and Drug Administration

GPI: glycosyl phosphatidylinositol

LDH: lactate dehydrogenase

PNH: paroxysmal nocturnal hemoglobinuria

STEC-HUS: Shiga toxin E. coli related

hemolytic uremic syndrome

TMA: thrombotic microangiopathy

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): patients with unresolved *Neisseria Meningitidis* infection; patients who are not currently vaccinated against *Neisseria meningitidis*, unless the risks of delaying Ultomiris treatment outweigh the risks of developing a meningococcal infection
- Boxed warning(s): serious meningococcal infections

Appendix D: General Information

- Ultomiris is only available through a REMS (Risk Evaluation and Mitigation Strategy) program due to the risk of life-threatening and fatal meningococcal infection. Patients should be vaccinated with a meningococcal vaccine at least 2 weeks prior to receiving the first dose of Ultomiris and revaccinated according to current medical guidelines for vaccine use. Patients should be monitored for early signs of meningococcal infections, evaluated immediately if infection is suspected, and treated with antibiotics if necessary.
- Examples of symptoms of anemia include but are not limited to: dizziness or lightheadedness, fatigue, pale or yellowish skin, shortness of breath, chest pain, cold hands and feet, and headache.

V. Dosage and Administration

Indication	Dosing Regimen*			Maximum Dose
PNH	Body Weight Range (kg)	Loading Dose (mg)	Maintenance Dose (mg)	3,600 mg/ 8 weeks
	≥ 40 to < 60	2,400	3,000 every 8 weeks	
	≥ 60 to < 100	2,700	3,300 every 8 weeks	
	≥ 100	3,000	3,600 every 8 weeks	
	Day 1: Loading dose IV Day 15 and thereafter: Maintenance dose IV			
aHUS	Body Weight Range (kg)	Loading Dose (mg)	Maintenance Dose (mg)	3,600 mg/ 8 weeks
	≥ 5 to < 10	600	300 every 4 weeks	
	≥ 10 to < 20	600	600 every 4 weeks	
	≥ 20 to < 30	900	2,100 every 8 weeks	
	≥ 30 to < 40	1,200	2,700 every 8 weeks	
	≥ 40 to < 60	2,400	3,000 every 8 weeks	
	≥ 60 to < 100	2,700	3,300 every 8 weeks	
	≥ 100	3,000	3,600 every 8 weeks	
	Day 1: Loading dose IV Day 15 and thereafter: Maintenance dose IV			

**For patients switching from eculizumab to Ultomiris, administer the loading dose of Ultomiris IV 2 weeks after the last eculizumab infusion, and then administer maintenance doses IV once at the specified frequency, starting 2 weeks after loading dose administration.*

VI. Product Availability

Single-dose vial: 300 mg/30 mL

VII. References

1. Ultomiris Prescribing Information. Boston, MA: Alexion Pharmaceuticals, Inc.; October 2019. Available at: www.ultomiris.com. Accessed October 22, 2019.
2. Parker C, Omine M, Richards S, et al. Diagnosis and management of paroxysmal nocturnal hemoglobinuria. Blood 2005; 106(12):3699-3709. doi:10.1182/blood-2005-04-1717.
3. Loirat C, Fakhouri F, Ariceta G, et al. An international consensus approach to the management of atypical hemolytic uremic syndrome in children. Pediatr Nephrol. 2016; 31: 15-39.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J1303	Injection, ravulizumab-cwvz, 300 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	02.19.19	05.19
1Q 2020 annual review: criteria added for new FDA indication: aHUS; references reviewed and updated.	11.26.19	02.20
WCG: adapting policy from CNC	12.8.20	12.20

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.